Mr. Gray on the Slender-tongued Saurians.

281

Inhab. Egypt.

Scapteira lineata. (Ida lineata, Gray, MSS. B. M.) Greenish; back and base of the tail with six bright blue longitudinal streaks; beneath yellowish; scales of the back rather large, rhombic, keeled; of the base of the tail short, sharply keeled.

Inhab. N. Africa.

b. Præanal shields similar, but narrower; abdominal shields 10 or 12-rowed, contracted behind; interparietal plate rudimentary.

Scapteira maculata. (Ida maculata, Gray, MSS. B. M.) Savigny Rept. Egypt, t. 1, f. 9.?? Greenish, with a broad pale dorsal streak, marked with indistinct blackish spots; sides of the back blackish, with three or four series of longitudinal oblong pale spots; scales of the back small, rhombic, acutely keeled; of the base of the tail broad, short; toes slightly fringed.

Inhab. Tripoli.

c. Præanal shields unequal; the hinder central one large, the rest smaller, placed in an arched series; abdominal plates, 12-rowed. Ida.

Scapteira inornata. (Ida inornata. Gray, MSS. B. M.) Greenish olive; tail paler, with a dark streak on each side; beneath silvery white; scales of the back small, ovate, acutely keeled; of the base of the tail broad; præanal scales 10 or 12, subequal, with small scales on the sides, placed in four transverse subalternating series.

Inhab. Tripoli.

Scapteira punctulata. (Ida punctulata, Gray, B. M.) Olive; back with six longitudinal series of small distant blackish dots; tail and cheeks brown varied; beneath whitish; scales of the back small, convex, rather rhombic, smooth; of the tail elongate, narrow.

Inhab. ——.

Scapteira nebulosa. (Ida nebulosa, Gray, MSS. B. M.) Pale olive; back with three series of rather large distant black spots; tail elongated, with a series of black dots on each side; scales of back minute, granular, smooth, flattish, of upper part of tail elongated, truncated.

Inhab. Egypt.

Scapteira leiocerca. (Ida leiocerca, Gray, MSS. B. M.) Olive black, varied with brown, leaving six series of small round pale spots, and six series of large oblong transverse pale spots, placed between the others; scales of the back small, rhombic, acute, smooth; of the base of the tail large, similar, smooth.

Inhab. ——.

The toes of S. inornata are longly, of S. inæqualis, S. maculata,