

Impact of the degree of urbanization on composition and structure of helminth communities in the Mongolian racerunner (*Eremias argus*) Peters, 1869

Z.N. Dugarov*, D.R. Baldanova and T.R. Khamnueva

Institute of General and Experimental Biology, Siberian branch, Russian Academy of Sciences, Ulan-Ude 670047, Russia

(Received 28 December 2016; Accepted 24 February 2017)

Abstract

The effects of the degree of urbanization on the composition and mean abundance of helminth species and the structure of helminth communities in the Mongolian racerunner were investigated along a rural–urban gradient in the region of Ulan-Ude city (the Republic of Buryatia, Russia) and neighbouring rural districts. Racerunners were obtained from key areas and categorized into three grades based on the degree of urbanization. In this study, a total of 208 lizards were examined. The helminth communities in the Mongolian racerunner were studied at the infracommunity and component community levels. The nematode *Spauligodon pseudoeremiasi* was a sensitive bioindicator of the degree of urbanization in our study. All parameters of helminth infracommunities in the Mongolian racerunner were significantly reduced with increasing degree of urbanization of the key areas. Two parameters of helminth component communities (the proportion of hosts infected with parasites and the Shannon index) were significantly reduced with increasing degree of urbanization. The decline recorded in parameters of helminth infracommunities and component communities in the Mongolian racerunner were probably connected with the attenuation of the relationships between helminths, having a complex life cycle with intermediate hosts along the rural–urban gradient.

Introduction

Cities with surrounding suburban areas represent a practical model for studying the effect of urbanization on discrete species and communities of species. McKinney (2008) showed that the species richness of plants, invertebrates, amphibians, reptiles and mammals decreases in the urban areas. The effect of a moderate degree of urbanization (suburban area) on the species richness of these groups of organisms is variable.

Urbanization affects both free-living and parasitic organisms. Growing cities increasingly intrude into the wild, and affect the distribution and transmission success of parasites. Continued rapid urbanization has increased

the importance of research on wildlife diseases in urban landscapes (Bradley & Altizer, 2007). Moreover, parasites are very sensitive indicators of environmental degradation (Vidal-Martinez & Wunderlich, 2017), caused by urbanization.

In developing countries, progressive urbanization is associated with the unplanned, uncontrolled and constant migration of people from rural to urban settlements. This migrant population brings, among other things, intestinal parasitic diseases, the level of which is already high in these settlements (Crompton & Savioli, 1993). The level of intestinal parasitoses in La Plata, Argentina was reduced as follows: a marginal zone > a suburban neighbourhood > an urban area (Gamboa *et al.*, 2003).

In a study on the impact of urbanization on the epidemiology of intestinal helminths in red foxes, in Geneva (Switzerland), it was shown that dioxenous

*E-mail: zhar-dug@biol.bscnet.ru

helminths that depend on rodent intermediate hosts are negatively affected by urbanization, as a result of the lower supply of rodents in the urbanized environments (Reperant *et al.*, 2007). Rodents may represent valuable indicators of infection pressure by free-living stages of *Echinococcus multilocularis*, *Toxocara* spp. and *Toxoplasma gondii* for humans in urban environments (Reperant *et al.*, 2009).

The presence of humans in cities and the associated altered urban landscape characteristics are associated with increased infections with gastrointestinal parasites in house finches (*Haemorhous mexicanus*) (Giraudeau *et al.*, 2014). Urbanization leads to a significant reduction in the helminth fauna of the Eurasian blackbird (*Turdus merula* L.), which is highly adapted to synanthropic habitats (Sitko & Zalesny, 2014). Environmental changes, such as changes in land use and wetland characteristics, nutrient runoff, pesticide contamination, biodiversity loss and climate change, have demonstrated the potential to enhance infection with *Ribeiroia* and *Echinostoma* in amphibians (Johnson & McKenzie, 2009). The above studies were related to the impact of urbanization on parasites of humans, mammals, birds and amphibians. Investigations of the effect of urbanization on reptilian parasites are still absent.

Reptiles perform important functions in natural ecosystems by serving as predators, prey for predators, pasture inhabitants, seed dispersers and synanthropes; they also serve as bioindicators of environmental conditions (Böhm *et al.*, 2013). The Mongolian racerunner (*Eremias argus*) is found in eastern Mongolia, China, the western part of the Korean Peninsula and Russia. In Russia, this racerunner is distributed in southern Buryatia and the south-western part of the Zabaikal'skiy region (Ananjeva *et al.*, 2004). The only helminthological study of the Mongolian racerunner within Zabaikalie was conducted by Sharpilo (1976). The aim of this research was to study the impact of the degree of urbanization on mean abundance, species composition and community structure of the parasites of *E. argus*.

Materials and methods

Samples

In this study, Mongolian racerunners were captured at five different sampling locations in the city of Ulan-Ude, and Ivolginsk and Tarbagatay regions adjacent to the city of Ulan-Ude. The five key areas are: (1) Ivolginsky Hollow (IH) (Ivolginsk District) (51°43'48"N, 107°22'10"E); (2) Upper Sayantuy (US) (Tarbagatay District) (51°42'22"N, 107°34'09"E); (3) Vakhmistrovo village (Va) (Tarbagatay District) (51°44'02"N, 107°33'14"E); (4) Silikatnyi village (Si) (Ulan-Ude city) (51°45'39"N, 107°33'05"E); (5) Neighbourhood No. 113 (Qu) (Ulan-Ude) (51°45'49"N, 107°35'52"E) (fig. 1).

The five key areas were categorized into three grades based on the degree of urbanization, in accordance with the classification of the degrees of urbanization (DEGURBA) of the European Commission (Dijkstra & Poelman, 2014), which is as follows: (1) thinly populated area (rural area) (outside urban clusters); (2) intermediate density area (towns and suburbs or small urban area) (a

density of at least 300 inhabitants/km²); (3) densely populated area (cities or large urban area) (a density of at least 1500 inhabitants/km²). These grades were applied for the statistical analysis of the data.

The key area IH refers to the thinly populated (rural) cluster (rank 1). It is located in the steppe area, outside human settlements. The locations within this key area where Mongolian racerunners may be obtained are scarcely susceptible to urbanization. They are separated by considerable distances from the populated areas, partly located within the forest shelter belts and partly within rocky areas with an extremely rugged terrain (these areas are not arable).

The key areas US and Va are moderately populated (suburbs – rank 2). The first of these areas is located near the eponymous village with a constant population of about 400 people. The area Va is close to the federal highway with heavy traffic. Several suburban settlements are located close to this key area.

The key areas Si and Qu refer to the cluster of densely populated areas (cities – rank 3). In the Si region, buildings and streets are seen. Apartment buildings are also seen in the second area, Qu.

In this study, Mongolian racerunners were obtained during the summers of 2009 and 2011–2012; numbers of specimens: IH, 39; US, 27; Va, 35; Si, 90 and Qu, 17. Most of the lizards (121 specimens) were accidentally caught in pitfall traps for soil insects and transferred to our laboratory for dissection. A total of 87 specimens were caught by hand. Captured lizards were sacrificed with a lethal injection of sodium thiopental. The lizards were dissected in the laboratory, post-mortem examinations of the body cavity, digestive tract, kidneys, liver, lungs, brain, eyes and muscles were examined for helminth parasites, using a stereomicroscope Stemi 2000 (Carl Zeiss, Göttingen, Germany). The worms found were fixed in 70% ethanol. Nematodes and acanthocephalans were mounted temporarily on slides and cleared with lactophenol or glycerin. Cestodes were stained with acetic carmine and, after dehydration, were mounted with Canada balsam. Worms were identified with the use of a microscope (AxioImager.M2, Carl Zeiss). All helminths were identified using the relevant keys (Petrochenko, 1958; Sharpilo, 1976). Worms were deposited in the helminthological collection of the Institute of General and Experimental Biology, Siberian branch, Russian Academy of Sciences.

Definitions

In this study, species richness was considered as the total number of parasite species found, and the mean species richness was the sum of the number of parasite species in each individual host divided by the number of hosts analysed. Mean number of specimens of helminth species was the sum of the number of helminth specimens in each individual host divided by the number of hosts analysed. Mean abundance was the total number of individuals of a particular parasite species in a sample of a particular host species divided by the total number of hosts of that species examined (including both infected and uninfected hosts). Prevalence was the number of hosts infected with one or more individuals of a particular




Fig. 1. Map of Baikal region, showing the localities where Mongolian racerunners were collected. Dark circles indicate sampling sites (IH, Ivolginsky Hollow; US, Upper Sayantuy; Va, Vakhmistrovo village; Si, Silikatnyi village; Qu, Neighbourhood No. 113). Polygons with grey squares indicate populated areas; 1, Selenga River; 2, road; 3, railway.

parasite species (or taxonomic group) divided by the number of hosts examined for that parasite species. The following definitions of parasite communities were applied: infracommunity – all parasite species within an individual host; component community – all parasite infracommunities within a host population (Bush & Holmes, 1986; Goater *et al.*, 1987; Bush *et al.*, 1997).

Evaluation of helminth communities

To evaluate the helminth infracommunities of the Mongolian racerunner, four parameters were applied: (1) species richness, (2) dominance, (3) evenness and (4) species diversity based on the relative abundance of

species. Species richness was evaluated based on the number of parasite species – mean species richness, Menhinick and Margalef indices; dominance – Berger–Parker index; evenness – Simpson, Camargo, Smith and Wilson evenness indices; species diversity was based on the relative abundance of species – Brillouin index. In addition to these indicators, the number of parasite specimens was determined. To characterize the helminth component communities, the total number of parasite species, the proportion of hosts infested with parasites, the Berger–Parker index of dominance and the Shannon index were used. These indices are widely used for studying various groups of organisms (Smith & Wilson, 1996; Magurran, 2004; Payne *et al.*, 2005).

Statistical analyses

The correlations between the mean abundances of the Mongolian racerunner helminths in various key areas and the degree of urbanization of these areas were calculated using Spearman's rank correlation coefficient. This analysis was repeated to establish the relationship between parameters of helminth communities of this host in various key areas and the degree of urbanization of these areas. Further, the mean abundance data of each helminth species of *E. argus* among key areas were compared using Mann-Whitney's *U*-test. This analysis was repeated to compare the parameters of helminth infracommunities of this host among the key areas. Thereafter, the proportions of *E. argus* specimens infected with parasites among the key areas were compared. Statistical analyses were carried out using STATISTICA 6.0 software (StatSoft Russia, Moscow, Russia).

Results

Fauna of helminths

Seven species of helminths were observed in the Mongolian racerunner: the cestodes *Oochoristica tuberculata* and *Mesocestoides lineatus*; the nematodes *Spauligodon pseudoeremiasi*, *Abbreviata abbreviata*, *Skrjabinelazia hoffmanni* and *Spirocerca lupi*; and the acanthocephalan *Macracanthorhynchus catulinus*. The composition of the helminth fauna of *E. argus* inhabiting discrete key areas ranged from two species (Qu) to six species (Va).

Two species of nematodes, *S. pseudoeremiasi* and *A. abbreviata*, were recorded in all key areas. The nematode *S. pseudoeremiasi* was predominant among helminths of the Mongolian racerunner in four key areas: IH, Va, US and Qu. Si was the only key area where *S. pseudoeremiasi* could not prevail. There was a statistically significant reduction in the mean abundance of *S. pseudoeremiasi* with increasing degree of urbanization, with its highest mean abundance being registered in the key area IH and the lowest in the key area Si. *Abbreviata abbreviata* was a subdominant species.

The high level of infection of *E. argus* with the cestode *O. tuberculata* was mainly observed in the key area IH, but rarely in the key area Va. *Oochoristica tuberculata* was not found in any of the other three areas. The cestode *M. lineatus* in the Mongolian racerunner was registered only in the key area Si, where it was found to be predominant. The nematode *S. hoffmanni* was found to have a low prevalence in the host in three key areas – Va, US and Si. Single specimens of acanthocephalan *M. catulinus* were observed in two key areas (IH and Va) and the nematode *S. lupi* in one (Va) (table 1).

The infracommunities of helminths

All indicators of the parasite infracommunities of *E. argus* (the number of species and number of specimens; species richness indices (Menhinick and Margalef); Berger-Parker index of dominance; evenness indices of Simpson, Camargo, Smith and Wilson; Brillouin diversity index) had negative, statistically significant correlations with the degree of urbanization (table 2).

The parameters of the parasite infracommunities of the Mongolian racerunner (the number of species and number of specimens; the indices of species richness of Menhinick and Margalef; the Berger-Parker index of dominance; Simpson, Camargo, and Smith and Wilson evenness indices; Brillouin diversity index) in the key area IH were statistically higher as compared with all the other key areas ($P < 0.01$).

The component communities of helminths

The proportion of host specimens infected with helminths in the component communities of *E. argus* decreased in a statistically significant manner with increasing degree of urbanization of the key areas. The Shannon index decreased in a statistically significant manner with increasing degree of urbanization of the key areas (table 3).

The proportion of the host specimens infected with helminths in the component communities of the Mongolian

Table 1. Correlations of mean abundances of helminths infecting Mongolian racerunner in different key areas with the degree of urbanization of these areas.

	Key areas					r_s	<i>P</i>
	IH	Va	US	Si	Qu		
Rank of the degree of urbanization	1	2	2	3	3		
Cestoda							
<i>Oochoristica tuberculata</i>	9.23	0.40	0	0	0	-0.825	0.086
<i>Mesocestoides lineatus</i>	0	0	0	6.28	0	0.559	0.327
Nematoda							
<i>Spauligodon pseudoeremiasi</i>	15.93	5.14	9.56	0.29	2.00	-0.949*	0.014
<i>Abbreviata abbreviata</i>	1.20	0.37	0.81	0.17	0.53	-0.738	0.155
<i>Skrjabinelazia hoffmanni</i>	0	0.11	1.07	0.27	0	0.135	0.828
<i>Spirocerca lupi</i>	0	0.06	0	0	0	-0.186	0.764
Acanthocephala							
<i>Macracanthorhynchus catulinus</i>	0.08	0.09	0	0	0	-0.648	0.237
The total number of parasite species	4	6	3	4	2		

*Statistically significant values of r_s .

Table 2. Correlations between parameters of Mongolian racerunner helminth infracommunities and the degree of urbanization of the key areas.

Parameter	Key areas					r_s	P
	IH	Va	US	Si	Qu		
Rank of the degree of urbanization	1	2	2	3	3		
Mean species richness	1.64	0.43	0.44	0.26	0.41	-0.498*	0.001
Mean number of specimens of helminth species	27.10	6.17	11.44	7.00	2.53	-0.507*	0.001
Menhinick index	0.41	0.21	0.17	0.15	0.17	-0.391*	0.001
Margalef index	0.25	0.04	0.01	0.01	0.08	-0.408*	0.001
Berger-Parker index	0.74	0.32	0.39	0.22	0.20	-0.391*	0.001
Dominant species	<i>S. p.</i>	<i>S. p.</i>	<i>S. p.</i>	<i>M. l.</i>	<i>S. p.</i>		
Simpson evenness index	0.76	0.24	0.33	0.12	0.14	-0.490*	0.001
Camargo evenness index	0.40	0.05	0.03	0.02	0.13	-0.407*	0.001
Smith and Wilson evenness index	0.72	0.33	0.41	0.21	0.17	-0.394*	0.001
Brillouin index	0.29	0.03	0.02	0.01	0.06	-0.418*	0.001
Number of examined Mongolian racerunner specimens	39	35	27	90	17		

*Statistically significant values of r_s . *S. p.*, *Spauligodon pseudoeremiasi*; *M. l.*, *Mesocestoides lineatus*.

racerunner in the key area IH was statistically higher as compared with all other key areas ($P < 0.001$).

Discussion

The nematode *S. pseudoeremiasi* was the dominant species among helminths of the Mongolian racerunner in four key areas. The life cycle of *S. pseudoeremiasi* is yet to be described. However, all nematodes of the order Oxyurida have a direct life cycle (Adamson, 1989). Nematodes with a direct life cycle, especially *Capillaria inaequalis* and *Thelandros magnavulvaris*, dominated the parasite fauna of the salamanders *Leurognathus marmorata*, *Desmognathus quadromaculatus*, *D. monticola* and *D. ochrophaeus* (Goater et al., 1987). It is possible that the direct life cycle of *S. pseudoeremiasi* may contribute to the dominance of the nematode among *E. argus* parasites. *Spauligodon pseudoeremiasi* was the only helminth of the lizard, whose infection reduced in a statistically significant manner with increasing degree of urbanization of key areas. Therefore, this parasite species is the most appropriate bioindicator (of the seven species studied) of the degree of urbanization in our research.

The life cycle of *A. abbreviata* is unknown. Studies have been carried out on *A. kazakhstanica*, whose intermediate hosts are 20 species of insects from five families

(Tenebrionidae, Tettigoniidae, Acrididae, Gryllidae and Manteidae) and three orders (Orthoptera, Coleoptera and Mantoptera). The diversity of the intermediate hosts contributes to the successful infection of the definitive hosts (Kabilov, 1980). The subdominant status of *A. abbreviata* in the Mongolian racerunner may testify to a wide range of its intermediate hosts. This, therefore, promotes the occurrence of this species in all key areas.

The cestode *O. tuberculata* was observed in *E. argus* with a high infection level only in the key area IH. It was found sporadically in the area Va, but not reported in the other three areas. The life cycle of *O. tuberculata* is yet to be identified. It has been found experimentally, in the North American species of the genus *Oochoristica*, that the life cycle of these cestodes is complex. Their intermediate hosts are beetles of the family Tenebrionidae and Dermestidae, larvae of butterflies of the family Pyralidae and grasshoppers of the family Acrididae (Rendtorff, 1948; Millemann, 1955; Widmer & Olsen, 1967; Conn, 1985; Criscione & Font, 2001). The high level of infection of the Mongolian racerunner with *O. tuberculata* in the key area IH can be attributed to the high level of trophic relations of lizards with intermediate hosts of the cestode, especially with darkling beetles, which are significantly abundant in this area.

Mesocestoides lineatus is the dominant species among helminths of the Mongolian racerunner only in the key

Table 3. Correlations between parameters of Mongolian racerunner helminth component communities and the degree of urbanization of the key areas.

Parameter	Key areas					r_s	P
	IH	Va	US	Si	Qu		
Rank of the degree of urbanization	1	2	2	3	3		
The total number of parasite species	4	6	3	4	2	-0.406	0.498
Proportion of host specimens infected with helminths	0.90	0.34	0.41	0.22	0.24	-0.949*	0.014
Berger-Parker index	0.60	0.83	0.84	0.90	0.79	0.527	0.361
Dominant species	<i>S. p.</i>	<i>S. p.</i>	<i>S. p.</i>	<i>M. l.</i>	<i>S. p.</i>		
Shannon index	0.81	0.68	0.56	0.35	0.51	-0.949*	0.014
Number of examined Mongolian racerunner specimens	39	35	27	90	17		

*Statistically significant values of r_s . *S. p.*, *Spauligodon pseudoeremiasi*; *M. l.*, *Mesocestoides lineatus*.

area Si. Data regarding the life cycle of species of the genus *Mesocestoides* are fragmented. This species has a complex life cycle. The first intermediate hosts of the genus *Mesocestoides* remain unidentified; it is only known that they are arthropods (insects, mites). The second intermediate hosts of *M. lineatus* are amphibians, reptiles, birds and small mammals; those in Buryatia are the brown rat *Rattus norvegicus*, the Chinese striped hamster *Cricetulus barabensis*, the house mouse *Mus musculus*, the Siberian chipmunk *Tamias sibiricus*, the long-tailed ground squirrel *Spermophilus undulatus* and the wood lemming *Myopus schisticolor*. The definitive hosts of *M. lineatus* are carnivorous mammals, mainly foxes, dogs and cats (Chertkova & Kosupko, 1978); those in Buryatia are the dog *Canis lupus familiaris*, the red fox *Vulpes vulpes*, the cat *Felis silvestris catus*, the Eurasian lynx *Lynx lynx*, the mountain weasel *Mustela altaica*, the sable *Martes zibellina* and the Pallas's cat *Felis manul* (Zhaltanova, 1992). The Mongolian racerunner is also a second intermediate host of *M. lineatus*. The infection of *E. argus* with tetrathyridia of *M. lineatus* in the key area Si only, indicates that there are already established stable trophic relations of definitive hosts of the cestode with the lizard in that area. Si is an industrial district of Ulan-Ude, with construction businesses and residential buildings. The presence of abandoned buildings in close proximity to apartment blocks contributes to an increase in the number of packs of stray dogs. Foxes have been observed in the area Si. It has been reported that foxes do not shy away from humans and have become synanthropic species in Europe (Scott *et al.*, 2014). *Mesocestoides lineatus* was the dominant species in this key area, instead of *S. pseudoeremiasi*.

Of the genus *Skrjabinelazia*, the life cycle of only *S. galliardi* has been studied previously. In *S. galliardi*, young females are larviparous, they produce thin-shelled eggs, infective stage 3 larvae and likely contribute to propagation within the host. Older large females produce numerous thick-shelled eggs containing infective stage 3 larvae. Eggs can survive in the ambient environment for at least 14 days. These resistant eggs ensure transmission to new hosts (Chabaud *et al.*, 1988). The life cycle of *S. hoffmanni* is probably similar to that of *S. galliardi*. The low level of infection of the Mongolian racerunner with *S. hoffmanni* is possibly related to the low degree of trophic relationships of the lizard with intermediate hosts (field crickets) within the area.

The life cycle of *M. catulinus* is complex; it includes intermediate, paratenic and definitive hosts. The intermediate hosts of this parasite are darkling beetles, *Adesmia gebleri* (Rizhikov & Dizer, 1954) and *Tentyria tessulata* (Farzaliev & Petrochenko, 1980). The definitive hosts of this parasite are carnivorous mammals (dog, fox, korsak, badger and others). *Macracanthorhynchus catulinus* is capable of maturing when transferred from one dog to another (Petrochenko, 1958; Sharpilo, 1976; Farzaliev & Petrochenko, 1980). The paratenic hosts of *M. catulinus* are represented by many species of amphibians, reptiles (including the Mongolian racerunner) and mammals.

The life cycle of the nematode *S. lupi* is complex, involving intermediate, paratenic and definitive hosts. The intermediate hosts of the nematode are various coprophagous beetles. The paratenic hosts of *S. lupi* are represented by

many species of reptiles, wild and domestic birds, as well as small mammals (rodents, rabbits and hedgehogs). The larvae of this nematode are transferred from one paratenic host to another. The final hosts are carnivores, mostly canids (dogs, wolves and foxes) (Bailey, 1972; Van der Merwe *et al.*, 2008).

The infection of definitive hosts (carnivores) with the acanthocephalan *M. catulinus* and the nematode *S. lupi* through intermediate hosts (beetles) is unlikely or extremely limited. The main source of infection of the definitive hosts with these two species of helminths is paratenic hosts, including reptiles (Sharpilo, 1983). Due to their low level of infection, they may be referred to as rare parasites of the Mongolian racerunner.

All indicators of the infracommunities (number of species and number of specimens; the indices of species richness by Menhinick and Margalef; Berger–Parker dominance index; indices of evenness by Simpson, Camargo, Smith and Wilson; Brillouin index), as well as the proportion of host specimens infected with parasites, showed a statistically significant decrease with increasing degree of urbanization (table 2). The decreased number of species and number of parasite specimens of *E. argus*, indices of species richness by Menhinick and Margalef, and the Brillouin index, as well as the proportion of infected host specimens in the helminth infracommunities with increasing degree of urbanization of the key areas may be attributed to the weakened relations between helminths with complex life cycles and the first intermediate hosts (insects). The decrease in the Berger–Parker dominance index in the helminth infracommunities was due to the diminished infection level of the lizard with the dominant species *S. pseudoeremiasi* with increasing degree of urbanization in the key areas.

In the component communities of helminths of *E. argus*, the proportion of host specimens infected with parasites decreased in a statistically significant manner with the increased degree of urbanization in the key areas. This is most likely due to the decreased diversity of insects, including intermediate hosts of helminths, with increase in the degree of urbanization. The statistically significant decrease of the Shannon index in the component communities of helminths of the Mongolian racerunner was based on the gradient of urbanization, reflecting the decrease in number of species and number of specimens in the infracommunities of helminths on this gradient.

Among the five key areas, IH was the only area which referred to the thinly populated (rural) cluster. The other areas were clusters with populations of intermediate and high density. The highest indicators of the infracommunities of parasites of the Mongolian racerunner (number of species and number of specimens, indices of species richness by Menhinick and Margalef; Berger–Parker index of dominance; evenness indices of Simpson, Camargo and Smith and Wilson; Brillouin index), as well as the proportion of host specimens infected with parasites were observed in key area IH. This area has the lowest degree of urbanization in all locations under study. The maximum level of infection with the nematode *S. pseudoeremiasi*, with most likely a direct life cycle, emphasizes the close relationships among specimens of *E. argus* within the key area. The highest levels of infection with the cestode *O. tuberculata* and the

nematode *A. abbreviata*, with complex life cycles, show stable trophic relationships between the lizard and intermediate hosts of helminths (insects). Maximum species richness (six species) was observed in the helminth component community of the Mongolian racerunner in the key area Va, which is due to host infection with *M. catulinus* and *S. lupi*. Both of these species are rare parasites of *E. argus*. At the same time, *S. lupi* was mainly observed in this key area. The infection of the Mongolian racerunner with *M. catulinus* and *S. lupi*, in which the definitive hosts are carnivorous mammals, typically canids, indicated strengthened trophic relationships of these mammals with the lizard. It is likely that the diversity of beetles (first intermediate hosts of these helminths) is relatively high in the Va key area. However, all indicators of the infracommunities and helminth component communities of this host in the key area Va, were noticeably less than in the IH area.

The impact of urbanization manifests through decreased species richness and relative abundance of helminths (Hartson *et al.*, 2011; Sitko & Zalesny, 2014; Calegario-Marques & Amato, 2014; Campiao *et al.*, 2017). However, an increase in helminth species richness has been observed in rodents inhabiting South-East Asia from forests towards agricultural areas and human settlements (Chaisiri *et al.*, 2010). The infection prevalence and magnitude of the multiple infections were greater for red colobus (*Procolobus rufomitratus*) in the fragmented than the unfragmented forest. An increase in species richness was recorded in parasite communities in the urban areas as compared to the natural areas, only in mammals. This may be due to parasites whose life cycles include humans and domestic animals (Gillespie & Chapman, 2008). The cestode *Rodentolepis straminea* was observed in wood mouse (*Apodemus sylvaticus*) only in urban woodland in the British Isles and not in rural and peri-urban areas (Rushworth *et al.*, 2016). In the present study, the cestode *M. lineatus* was found in the Mongolian lizard only in the urban area and was not found in the rural area and suburbs. *Eremias argus* was involved in the circulation of *M. lineatus*, whose final hosts are canids (dogs and foxes) associated with humans, only in the urban area.

In conclusion: urbanization causes a significant change in the structure of helminth infracommunities of the Mongolian racerunner. All indicators of the parasite infracommunities of *E. argus* (the number of species and number of specimens; species richness indices (Menhinick and Margalef); Berger–Parker index of dominance; evenness indices of Simpson, Camargo, Smith and Wilson; Brillouin diversity index) and the proportion of host specimens infected with helminths were decreased with increasing degree of urbanization.

Spauligodon pseudoeremiasi can be a sensitive bioindicator of the degree of urbanization, because it is a dominant parasite of the Mongolian lizard and its relative abundance decreases most significantly with increasing degree of urbanization.

There are complex interactions between wildlife, domestic animals and humans in urban areas. Parasites infecting humans are involved in the processes of these interactions. Increasing attention of researchers is necessary for the study of changes in the species composition and structure of parasite communities of reptiles on a

gradient of urbanization. There is no doubt that in the course of these studies the role of reptiles in the transmission of zoonotic diseases will be more clearly identified.

Acknowledgements

The authors sincerely thank Natalia Schepina from the Geological Institute of the Siberian Branch of the RAS (Russian Academy of Sciences) and Larisa Khobrakova from the Institute of General and Experimental Biology of the Siberian Branch of the RAS for their help in obtaining the Mongolian racerunner specimens.

Financial support

This research work was carried out within the theme ‘Ecology of parasites of animals in the ecosystems of Baikal region: spatial distribution and parasite–host relationships’ (registration no. AAAA-A17-117011810039-4).

Conflict of interest

None.

Ethical standards

All animal procedures were conducted in accordance with the ethical standards established by the Russian national instructions for the care and use of laboratory animals. There is a Commission on Bioethics in the Institute of General and Experimental Biology of SB RAS, which examines all research projects with the use of animals.

References

- Adamson, M.L. (1989) Evolutionary biology of the Oxyurida (Nematoda): biofacies of a haplodiploid taxon. *Advances in Parasitology* **28**, 175–228.
- Ananjeva, N.B., Orlov, N.L., Khalikov, R.G., Darevsky, I.S., Ryabov, S.A. & Barabanov, A.V. (2004) *Colored atlas of the reptiles of the North Eurasia (taxonomic diversity, distribution, conservation status)*. St. Petersburg, Zoological Institute of the Russian Academy of Sciences (in Russian).
- Bailey, W.S. (1972) *Spirocerca lupi*: a continuing inquiry. *Journal of Parasitology* **58**, 3–22.
- Böhm, M., Collen, B., Baillie, J.E.M. *et al.* (2013) The conservation status of the world’s reptiles. *Biological Conservation* **157**, 372–385.
- Bradley, C.A. & Altizer, S. (2007) Urbanization and the ecology of wildlife diseases. *Trends in Ecology and Evolution* **22**, 95–102.
- Bush, A.O. & Holmes, J.C. (1986) Intestinal helminths of lesser scaup ducks: an interactive community. *Canadian Journal of Zoology* **64**, 142–152.
- Bush, A.O., Lafferty, K.D., Lotz, J.M. & Shostak, A.W. (1997) Parasitology meets ecology on its own terms: Margolis *et al.* revisited. *Journal of Parasitology* **83**, 575–583.
- Calegario-Marques, C. & Amato, S.B. (2014) Urbanization breaks up host–parasite interactions: a case study on

- parasite community ecology of rufous-bellied thrushes (*Turdus rufiventris*) along a rural–urban gradient. *PLoS ONE* **9**, e103144.
- Campiao, K.M., Ribas, A.C.A., Silva, I.C.O., Dalazen, G. T. & Tavares, L.E.R.** (2017) Anuran helminth communities from contrasting nature reserve and pasture sites in the Pantanal wetland, Brazil. *Journal of Helminthology* **91**, 91–96.
- Chabaud, A.G., Bain, O. & Poinar, G.O.** (1988) *Skrjabinelazia galliardi* (Nematoda, Securotoidea): complements morphologiques et cycle biologique. *Annales de Parasitologie Humaine et Comparee* **63**, 278–284.
- Chaisiri, K., Chaeychomsri, W., Siruntawineti, J., Bordes, F., Herbreteau, V. & Morand, S.** (2010) Human-dominated habitats and helminth parasitism in Southeast Asian murids. *Parasitology Research* **107**, 931–937.
- Chertkova, A.N. & Kosupko, G.A.** (1978) Suborder Mesocostoidata Skrjabin, 1940. pp. 118–129 in Ryzhikov, K.M. (Ed.) *Principles of cestodology, vol. 9*. Moscow, Nauka (in Russian).
- Conn, D.B.** (1985) Life cycle and postembryonic development of *Oochoristica anolis* (Cyclophyllidea: Linstowiidae). *Journal of Parasitology* **71**, 10–16.
- Criscione, C.D. & Font, W.F.** (2001) Development and specificity of *Oochoristica javaensis* (Eucestoda: Cyclophyllidea: Anoplocephalidae: Linstowiinae). *Comparative Parasitology* **68**, 149–155.
- Crompton, D.W.T. & Savioli, L.** (1993) Intestinal parasitic infections and urbanization. *Bulletin of the World Health Organization* **71**, 1–7.
- Dijkstra, L. & Poelman, H.** (2014) A harmonised definition of cities and rural areas: The new degree of urbanisation. Regional working Paper, 01/2014. Directorate-General for Regional and Urban Policy. Available at http://ec.europa.eu/regional_policy/sources/docgener/work/2014_01_new_urban.pdf (accessed 10 March 2017).
- Farzaliev, A.M. & Petrochenko, V.I.** (1980) New data on the life cycle of the acanthocephalan *Macracanthorhynchus catulinus* Kostylew, 1927 (Acanthocephala), a parasite of carnivores. *Trudy Vsesoyuznogo Instituta Gelmintologii* **25**, 140–144 (in Russian).
- Gamboa, M.I., Basualdo, J.A., Cordoba, M.A., Pezzani, B.C., Minvielle, M.C. & Lahitte, H.B.** (2003) Distribution of intestinal parasitoses in relation to environmental and sociocultural parameters in La Plata, Argentina. *Journal of Helminthology* **77**, 15–20.
- Gillespie, T.R. & Chapman, C.A.** (2008) Forest fragmentation, the decline of an endangered primate and changes in host parasite interactions relative to an unfragmented forest. *American Journal of Primatology* **70**, 222–230.
- Giraudeau, M., Mousel, M., Earl, S. & McGraw, K.** (2014) Parasites in the city: degree of urbanization predicts poxvirus and coccidian infections in house finches (*Haemorhous mexicanus*). *PLoS ONE* **9**, e86747.
- Goater, T.M., Esch, G.W. & Bush, A.O.** (1987) Helminth parasites of sympatric salamanders: ecological concepts at infracommunity, component and compound community levels. *The American Midland Naturalist* **118**, 289–300.
- Hartson, R.B., Orlofske, S.A., Melin, V.E., Dillon, R.T. & Johnson, P.T.J.** (2011) Land use and wetland spatial position jointly determine amphibian parasite communities. *EcoHealth* **8**, 485–500.
- Johnson, P.T.J. & McKenzie, V.J.** (2009) Effects of environmental change on helminth infections in amphibians: exploring the emergence of *Ribeiroia* and *Echinostoma* infections in North America. pp. 249–280 in Fried, B. & Toledo, R. (Eds) *The biology of echinostomes: from the molecule to the community*. New York, Springer.
- Kabilov, T.K.** (1980) Life cycle of the nematode, *Abbreviata kazachstanica*. *Parazitologiya* **14**, 263–270 (in Russian with an English summary).
- Magurran, A.E.** (2004) *Measuring biological diversity*. Oxford, Blackwell.
- McKinney, M.L.** (2008) Effects of urbanization on species richness: a review of plants and animals. *Urban Ecosystems* **11**, 161–176.
- Millemann, R.E.** (1955) Studies on the life-history and biology of *Oochoristica deserti* n. sp. (Cestoda: Linstowiidae) from desert rodents. *Journal of Parasitology* **41**, 424–440.
- Payne, L.X., Schindler, D.E., Parrish, J.K. & Temple, S.A.** (2005) Quantifying spatial pattern with evenness indices. *Ecological Applications* **15**, 507–520.
- Petrochenko, V.I.** (1958) *Acanthocephala of domestic and wild animals, vol. 2*. Moscow, Izdatelstvo Akademii Nauk SSSR (in Russian).
- Rendtorff, R.C.** (1948) Investigations on the life cycle of *Oochoristica ratti*, a cestode from rats and mice. *Journal of Parasitology* **34**, 243–252.
- Reperant, L.A., Hegglin, D., Fischer, C., Kohler, L., Weber, J.-M. & Deplazes, P.** (2007) Influence of urbanization on the epidemiology of intestinal helminths of the red fox (*Vulpes vulpes*) in Geneva, Switzerland. *Parasitology Research* **101**, 605–611.
- Reperant, L.A., Hegglin, D., Tanner, I., Fischer, C. & Deplazes, P.** (2009) Rodents as shared indicators for zoonotic parasites of carnivores in urban environments. *Parasitology* **136**, 329–337.
- Rizhikov, K.M. & Dizer, Y.B.** (1954) Biology of *Macracanthorhynchus catulinus* and *Mediorhynchus micracanthus*. *Doklady Akademii Nauk SSSR* **95**, 1367–1369 (in Russian).
- Rushworth, R.L., Boufana, B., Hall, J.L., Brannan, V., Mastin, A., Birtles, R.J., Craig, P.S. & Rogan, M.T.** (2016) *Rodentolepis straminea* (Cestoda: Hymenolepididae) in an urban population of *Apodemus sylvaticus* in the UK. *Journal of Helminthology* **90**, 476–482.
- Scott, D.M., Berg, M.J., Tolhurst, B.A., Chauvenet, A.L. M., Smith, G.C., Neaves, K., Lochhead, J. & Baker, P.J.** (2014) Changes in the distribution of red foxes (*Vulpes vulpes*) in urban areas in Great Britain: findings and limitations of a media-driven nationwide survey. *PLoS ONE* **9**, e99059.
- Sharpilo, V.P.** (1976) *Parasitic worms of the reptiles of the fauna of the USSR*. Kiev, Naukova dumka (in Russian).
- Sharpilo, V.P.** (1983) Reptiles of the fauna of the USSR, intermediate and reservoir hosts of helminths. *Parazitologiya* **16**, 177–184 (in Russian with an English summary).
- Sitko, J. & Zalesny, G.** (2014) The effect of urbanization on helminth communities in the Eurasian blackbird (*Turdus merula* L.) from the eastern part of the Czech Republic. *Journal of Helminthology* **88**, 97–104.

- Smith, B. & Wilson, J.B.** (1996) A consumer's guide to evenness indices. *Oikos* **76**, 70–82.
- Van der Merwe, L.L., Kirberger, R.M., Clift, S., Williams, M., Keller, N. & Naidoo, V.** (2008) *Spirocerca lupi* infection in the dog: a review. *Veterinary Journal* **176**, 294–309.
- Vidal-Martinez, V.M. & Wunderlich, A.C.** (2017) Parasites as bioindicators of environmental degradation in Latin America: a meta-analysis. *Journal of Helminthology* **91**, 165–173.
- Widmer, E.A. & Olsen, O.W.** (1967) The life history of *Oochoristica osheroffi* Meggitt, 1934 (Cyclophyllidea: Anoplocephalidae). *Journal of Parasitology* **53**, 343–349.
- Zhaltsanova, D.-S.D.** (1992) *Helminths of mammals of the Lake Baikal basin*. Moscow, Nauka (in Russian).