

HELMINTHOLOGIA, 54, 2: 174 - 178, 2017

Research Note

The first helminth study on Brandt's Persian Lizard *Iranolacerta brandtii* (De Filippi, 1863) (Squamata: Lacertidae) from Van Province, TurkeyS. BIRLIK^{1*}, H. S. YILDIRIMHAN¹, Y. KUMLUTAŞ², K. CANDAN², Ç. ILGAZ²

¹Uludag University Faculty of Arts and Science, Department of Biology, 16059 Nilüfer-Bursa, Turkey, E-mail: *sezen@uludag.edu.tr, yhikmet@uludag.edu.tr; ²Dokuz Eylül University, Faculty of Science, Department of Biology, 35160, Buca-Izmir, Turkey, E-mail: yusuf.kumlutas@deu.edu.tr, cetin.ilgaz@deu.edu.tr, kamilcandan@yahoo.com

Article info

Received January 23, 2017
Accepted March 10, 2017

Summary

Population of *Iranolacerta brandtii* lizard including 56 (21♀, 34♂, 1 juvenile) samples were examined for helminths. 17 of them were found to harbour one or more helminth parasites, while 39 didn't harbor any. Two helminth species were determined. These belong to Phylum: Nematoda, *Spauligodon aloisei* Casanova, Milazzo, Ribas & Cagnin, 2003 and *Skrjabinodon medinae*. García-Calvente, 1948. This study represents new host and locality records in both host lizard and the helminth species. *Sp. aloisei* is recorded for the first time; *Sk. medinae* is the third report from Turkey nevertheless *Sk. medinae* is a new host record for *I. brandtii*.

Keywords: *I. brandtii*; Helminth; Nematoda; Van; Turkey

Introduction

Parasites may account for half of the total number of species in the world but only a fraction of the diversity within these organisms has been detected and fewer described. (Maia, 2015). Reptiles are commonly infected by a wide range of parasites, serving either as their definitive or intermediate hosts (Al-Moussawi, 2010). Although Turkey has rich reptile fauna in the world, studies on the helminth fauna in these hosts are very scarce.

Lacertidae family contains approximately 39 species in Turkey (Baran & Atatür, 1998; Sindaco *et al.*, 2000; Arnold *et al.*, 2007; Baran *et al.*, 2012). Some helminth studies have been carried out on lizard species including in this genus in Turkey. These species are European Green Lizard, *Lacerta viridis* (Laurenti, 1768) (Schad *et al.*, 1960); Crimean Wall Lizard *Podarcis tauricus* (Pallas, 1814) (Schad *et al.*, 1960); Dwarf Lizard, *Parvilacerta parva* (Boulenger, 1887) (Saygı & Olgun, 1993); Danford's Lizard, *Anatololacerta danfordi* (Günther, 1876) (Gürelli *et al.*, 2007), Balkan Emerald Lizard, *L. trilineata* Bedriaga, 1886 (Yıldırımhan *et al.*, 2011); Pleske's Racerunner-Transcaucasian Racerunner, *Eremias pleskei*

Nikolsky, 1905 Strauch's Racerunner, *E. strauchi* Kessler, 1878, Suphan Racerunner, *E. suphani* Başoğlu & Helmich, 1968 (Düsen *et al.*, 2013) Anatolian Lizard, *Apathya cappadocica* (Werner, 1902) (Birlik *et al.*, 2015), Spiny-Tailed Lizard, *Darevskia rudis* (Bedriaga, 1886) (Roca *et al.*, 2015a) *D. uzzelli* (Darevsky, 1977), *D. bendimahiensis* (Schmidtler, Eiselt & Darevsky, 1994), *D. saphirina* (Roca *et al.*, 2015b) (Schmidtler, Eiselt & Darevsky, 1994), *D. clarkorum* (Darevsky & Vedmederja, 1977) *D. raddei* (Boettger, 1892), *D. parvula* (Lantz & Cyrén, 1913), *D. valentini* (Boettger, 1892), *D. armeniaca* (Méhely, 1909) *D. unisexualis* (Darevsky, 1966) (Roca *et al.*, 2016); Lebanon lizard *Phoenicolacerta laevis* (Gray, 1838) (Birlik *et al.*, 2016), *Acanthodactylus harranensis* Baran, Kumlutaş, Lanza, Sindaco, Ilgaz, Avci & Crucitti, 2005 *A. schreiberi* Boettger, 1878 *Mesalina brevirostris* Blanford, 1874 (Düsen *et al.*, 2016). *Iranolacerta brandtii* (De Filippi, 1863) has been reported for the first time from Van province in eastern Anatolia neighboring to Turkish and Iranian border (Avci *et al.*, 2015). The aim of this study is to evaluate the helminth fauna of *I. brandtii* from Turkey.

* – corresponding author




Fig.1. *Spauligodon aloisei* female anterior (prebulbar vulva) 10×.




Fig.2. *Spauligodon aloisei* female posterior 10×.

Table 1. The prevalence, mean intensity, abundance and numbers of minimum/maximum helminth and infected host.

Species of helminth	Number of host infected	Helminthes (minimum/maximum)	Prevalence (%)	Mean Intensity	Abundance
<i>Spauligodon aloisei</i>	11	1 – 10	19	2.72	0.53
<i>Skrjabinodon medinae</i>	14	1 – 6	25	2.85	0.71

Materials and Methods

Fifty-six specimens of *I. brandtii* were collected from two populations (Karadulda Village, Çaldıran and Saray, Van province, eastern Turkey) between July in 2015 and 2016. The snout vent length (SVL) of lizard samples was measured with a digital caliper to the nearest 0.01mm.

The body cavity was opened, and the digestive tract removed. The oesophagus, stomach, small and large intestine and lungs were opened and examined for helminths under dissecting microscope. Nematodes were killed in hot saline solution, fixed in 70 % ethanol, and mounted in glycerol. For the morphological examination, the helminth species were cleared gradually in glycerin. Parasites were identified to species level and the number and location of individuals of each species were recorded. Helminth identification was based on keys given by Yamaguti (1961), Petter and Quentin (1976), Baker (1987), Anderson (2000), Anderson and Chabaud (2009). Helminth voucher specimens were deposited in the Department of Biology, Uludağ University, Bursa, Turkey; lizard specimens were deposited in the Department of Biology, Dokuz Eylül University, İzmir, Turkey.

Results

Mean snout-vent length of lizard specimens was found as 68.57 ± 11.46 mm with a range from 46.0 to 87.0 mm. During

a helminthological examination of lacertid lizards from Turkey, 17 of 56 *I. brandtii* specimens were found to harbor 70 (9 males and 61 females) nematodes of the *Spauligodon* and the *Skrjabinodon*. The total infection rate of lizards by helminths was 30 %, with 17 of 56. These animals were host to 2 species of Nematoda. Total parasite number was 70. Seven of the 56 lizard specimens (12 %) were infected by two nematode species while 10 (17 %) were infected by one nematode species. The prevalence, mean intensity, abundance and other numerical data are presented in Table 1.



Fig.3. *Spauligodon aloisei* male posterior 10×.



Fig.4. *Skrjabinodon medinae* female anterior post bulbar vulva 10x.

Discussion

Three genera of pharyngodonid nematodes are commonly found in reptiles and exhibit a vulvar opening in the anterior part of the body, just posterior to the postbulbar excretory pore: Two of these genera are *Skrjabinodon* Inglis, 1968 and *Spauligodon* Skrjabin, Schikhobalova and Lagodovskaja, 1960. These genera can be separated by the relationship of the caudal alae to the genital papillae (Bursey & Goldberg, 1995).

Pharyngodonid nematodes of *Spauligodon* currently including 50 species are parasites of reptiles. Based upon the zoogeographic regions described by Holt *et al.* (2013) 16 *Spauligodon* species is known from the Palaearctic region. The Turkish helminth fauna represents 5 species including the *Spauligodon*. These species are *Spauligodon* sp. from *Parvilacerta parva* (Saygi & Olgun 1993), *Sp. atlanticus*, Astasio-Arbiza *et al.*, 1987 from *Apathya cappadocica* (Birlik *et al.*, 2015), *Sp. eremiasi* Markov & Bogdanov, 1961 from *Eremias trauchi* and *E. suphani* (Düşen *et al.*, 2013); *Sp. laevicauda* Seurat, 1924 from *Hemidactylus turcicus*; *Sp. saxicola*, Sharpilo, 1961 from *Eremias trauchi*, *E. suphani* (Düşen *et al.*, 2013), *Darevskia clarkorum*, *D. parvula*, *D. raddei*, *D. valentini*, *D. unisexualis* (Roca *et al.*, 2016).

Spauligodon aloisei was described in Italy (Casanova *et al.*, 2003). It was described as parasite of *Podarcis siculus* (Reptilia: Lacertidae). In Turkey, this helminth species is reported for the first time (Figs. 1, 2, 3). While spines are found in the tail filament of male specimens, there is no spicule that is diagnostic characters of the *Spauligodon*. The male specimens also have lateral alae. The female specimens found in lizard specimens having prebulbar vulva position that the most important diagnostic character for *S. aloisei*. *S. aloisei* is only one species having prebulbar vulva location for females in Palearctic region. All of the morphological and metrical features of our specimens were in agreement with the information given in the previous literature for *S. aloisei*.

Two species of *Skrjabinodon* Inglis, 1968 are currently known from

Turkey. These species are *Skrjabinodon aegyptiacus* Moravec, Barus & Rysavy, 1987 and *Sk. medinae*, García-Calvente, 1948. *Sk. medinae* was reported from *L. trilineata* (Yildirimhan *et al.* 2011), *A. cappadocica* (Birlik *et al.*, 2015) and *P. laevis* (Birlik *et al.*, 2016). *Sk. aegyptiacus* was reported from *Chalcides ocellatus* (Forsk., 1775) (Incedogan *et al.*, 2014). Males of *Skrjabinodon* do not have caudal alae and all genital papillae are embedded in the caudal alae (Figs. 4 and 5). Our male specimens have a spicule and the length of the spicule ranges from 55 to 60 μ . Obtained data are compatible with the data given in the previous literature for *Sk. medinae*. The female specimens have postbulbar vulva position (Fig. 4). They also possess a single pair of sessile pre-clacal papillae. In this study, *Sk. medinae* is a new host record for *I. brandtii*.

In our study, *Spauligodon* and *Skrjabinodon* have been found in coexistence. This correlates with usual pattern found in different European lacertid lizards, in which other *Spauligodon* species frequently share reptile hosts with members of the genera *Skrjabinodon* (Jorge *et al.*, 2014).

The component community of the helminths in amphibians and reptiles present fundamental differences in having low parasite richness when compared with other classes of vertebrates. Because they are poikilothermic animals, the environmental condi-




Fig.5 *Skrjabinodon medinae* male posterior (spicule) 40x.

tions limit the recruitment of potential parasite helminths, and the development of the parasitic community is affected by the feeding rates, as well as the forage behavior. (Aho, 1990) Besides all this, low infection rates and low species richness are correlated with a strictly entomophagous diet in many studies (Roca *et al.*, 1989; Galdón *et al.*, 2006; Roca *et al.*, 2006).

Until this time, there have been many studies on the helminth fauna of reptile species in Turkey. In these studies, many helminth species have been identified and it has been determined that the nematodes are dominant helminth communities in lacertid hosts. However, these studies are still inadequate to evaluate helminth fauna of Turkey. To identify new species, contributing biodiversity of Turkey, linking host characteristics with their parasite helminthes, further researches on helminth fauna of this group are needed.

References

- AHO, J.M. (1990): Helminth communities of amphibians and reptiles: Comparative approaches to understanding patterns and processes. In: ESCH, G.W., BUSH, A.O., AHO, J.M. (Eds) *Parasite communities: Patterns and processes*. Chapman and Hall. London, U.K. pp. 157 – 195
- ANDERSON, R.C. (2000): *Nematoda parasites of vertebrates, their development and transmission*. 2nd Edition, Walingford, UK, CABI Publishing, 650 pp.
- ANDERSON, R., CHABAUD, S. (2009): *Keys to the Nematode Parasites of Vertebrates*. Archival Volume, 480 pp.
- AVCI, A., ILGAZ, C. BOZKURT, E., UZUM, N., OLGUN, K. (2015): A new record of *Iranolacerta brandtii* (De Filippi, 1863) (Sauria: Lacertidae) in Eastern Anatolia, Turkey. *Russian J Herpetol.*, 22(1): 68 – 74
- AL-MOUSSAWI, A.A. (2010): First record in Iraq of *Tanqua anomala* (Linstow, 1904) from the dice snake, *Natrix tessellata tessellata* (Laurenti, 1768). *Bull. Iraq Nat. Hist. Mus.*, 11(1): 27 – 38
- ARNOLD, E.N., ARRIBAS, O., CARRANZA, A.S. (2007): Systematics of the palaeartic and oriental lizard tribe Lacertini (Squamata: Lacertidae: Lacertinae), with description of eight new genera. *Zootaxa*, 1430(1): 1 – 86. DOI: 10.11646/zootaxa.1430.1.1
- BAKER, M.R. (1987): Synopsis of the Nematoda Parasitic in Amphibians and Reptiles. *Memorial University of Newfoundland, Occasional Papers in Biology*. St. John's Newfoundland, Canada, 11: 1 – 325
- BARAN, I., ATATUR, M.K. (1998): *Türkiye Herpetofaunası (Kurbağa ve Sürüngenler) [Herpetofauna of Turkey (Amphibia and Reptilia)]*, Ankara (Çevre Bakanlığı), 214 pp. (In Turkish).
- BARAN, I., ILGAZ, C, AVCI, A., KUMLUTAS, Y., OLGUN, K. (2012): *Amphibians and Reptiles of Turkey*. Ankara (TÜBİTAK) 204 pp. (In Turkish)
- BIRLIK, S., YILDIRIMHAN, H.S., SUMER, N., KUMLUTAS, Y., ILGAZ, C., GUCLU, O., DURMUS, S.H. (2015): The helminth fauna of *Apathya cappadocica* (Werner, 1902) (Anatolian Lizard) from Turkey. *Helminthologia*, 52(4): 310 – 315. DOI: 10.1515/helmin-2015-0049.
- BIRLIK, S., YILDIRIMHAN, H.S., SUMER, N., KUMLUTAS, Y., ILGAZ, C., DURMUS, S.H., GUCLU, O., CANDAN, K. (2016): Helminth fauna of Lebanon Lizard, *Phoenicolacerta laevis* (Gray, 1838), (Squamata: Lacertidae) from Southern Turkey. *Helminthologia*, 53(3): 262 – 269. DOI: 10.1515/helmin-2016-0016
- BURSEY, C.R., GOLDBERG, S.R. (1995): *Spauligodon caymanensis* sp. n. (Nematoda: Pharyngodonidae) from *Anolis conspersus* (Sauria: Polychridae) from Grand Cayman Island, British West Indies. *J. Helminthol. Soc. Wash.*, 62(2): 183 – 187
- CASANOVA, J.C., MILAZZO, C., RIBAS, A., CAGNIN, M. (2003): *Spauligodon aloisei* n. sp. (Nematoda: Pharyngodonidae) parasite of *Podarcis sicula* (Reptilia: Lacertidae) from Italy. *J. Parasitol.*, 89(3): 577 – 579. DOI: 10.1645/0022-3395(2003)089[0577:SANSNP]2.0.CO;2
- DUSEN, S., KUMLUTAS, Y., ILGAZ, C., YAKA, H., KARADAYI, F. (2013): Helminth Parasites of the Three Racerunner Lizards: *Eremias pleskei* Nikolsky, 1905 (Pleske's Racerunner-Transcaucasian Racerunner), *Eremias trauchi* Kessler, 1878 (Strauch's Racerunner) and *Eremias suphani* Basoglu & Hellmich, 1968 (Suphan Racerunner) collected from Eastern Part of Turkey. *Helminthologia*, 50(2): 108 – 111, 2013. DOI 10.2478/s11687-013-0117-3
- DUSEN, S., KUMLUTAS, Y., ILGAZ, C., YAKAGUL, C. (2016): A helminthological research on three Lacertid lizards species: *Acanthodactylus harranensis* Baran *et al.*, 2005, *Acanthodactylus schreiberi* Boulenger, 1878, and *Mesalina brevisrostris* Blanford, 1874, collected from South and South-eastern regions of Turkey. *Helminthologia*. 53(2). DOI: 10.1515/helmin-2016-0010
- GALDON, M.A., ROCA, V., BARBOSA, D., CARRETERO, M.A. (2006): Intestinal helminth communities of *Podarcis bocagei* and *Podarcis carbonelli* (Sauria: Lacertidae) in NW Portugal. *Helminthologia*, 43: 37 – 41. DOI: 10.2478/s11687-006-0008-y
- GURELLI, G., GOCMEN, B., CETIN-DOGAN, T., ALPAGUT-KESKIN, N. (2007): First record of *Mesocestoides* spp. Vaillant, 1863 Tetrathyridia (Cestoidea: Cyclophyllidae) in Anatolian lizard, *Anatololacerta danfordi* (Günther, 1876) in Turkey. *North-Western J. Zool.*, 3(2): 96 – 104
- HOLT, B.G., LESSARD, M.K., BORREGAARD, S.A., FRITZ, M.B., ARAUJO, D., FABRE, P.H., GRAHAM, C.H., GAVES, G.R., JONSSON, K.A. (2013): An update of Wallace's zoogeographic regions of the World. *Science*, 339: 74 – 78. DOI: 10.1126/science.1228282
- INCEDOGAN, S., YILDIRIMHAN, H.S., BURSEY, C.R. (2014): Helminth parasites of the ocellated skink, *Chalcides ocellatus* (Forskal, 1775) (Scincidae) from Turkey. *Comp. Parasitol.*, 81(2): 260 – 269. DOI: 10.1654/4708.1
- JORGE, F., PERERA, A., ROCA, V., CARRETERO, M.A., HARRIS, D.J., POULIN, R. (2014): Evolution of alternative male morphotypes in oxyurid nematodes: a case of convergence? *J. Evol. Biol.*, 27: 1631 – 1643. DOI: 10.1111/jeb.12430
- MAIA, J. (2015): *Diversity, infection patterns and hostparasite associations of apicomplexan parasites in reptiles*. Tese de Doutorado apresentada à Faculdade de Ciências da Universidade do Porto Biologia

- PETTER A.J., QUENTIN J.C. (1976): *CIH keys to the nematode parasites of vertebrates*. No. 4 keys to genera of the Oxyuroidea, 1st ed. Commonwealth Agricultural Bureaux International, Farnham Royal, U.K., 30 pp.
- ROCA, V., LOPEZ-BALAGUER, E., HORNERO, M.J. (1989): Helminthofauna de *Podarcis hispanica* (Steindachner, 1870) y *Podarcis bocagei* (Seoane, 1884) (Reptilia: Lacertidae) en el Cuadrante Noroccidental de la Península Ibérica [Helminthofauna from *Podarcis hispanica* (Steindachner, 1870) and *Podarcis bocagei* (Seoane, 1884) (Reptilia: Lacertidae) in the North-West of the Iberian Peninsula]. *Rev. Ibér. Parasitol.*, 49: 127 – 135 (In Spanish)
- ROCA, V., CARRETERO, M.A., MARQUES, A., BARBOSA, D., GALDON, M.A. (2006): Relationships between helminth communities and host traits in *Podarcis bocagei* and *P. carbonelli* from NW Portugal. *Ital. J. Zool.*, 73(3): 213 – 217. DOI: 10.1080/11250000600727634.
- ROCA, V., JORGE, F., ILGAZ, C., KUMLUTAS, Y., DURMUS, S.H., CARRETERO, M.A. (2015a): The intestinal helminth community of the spiny-tailed lizard *Darevskia rudis* (Squamata, Lacertidae) from Northern Turkey. *J. Helminthol.*, 90(2): 144 – 151. DOI: 10.1017/S0022149X14000911
- ROCA, V., JORGE, F., ILGAZ, C., KUMLUTAS, Y., DURMUS, S.H., CARRETERO, M.A. (2015b): Are the helminth communities from unisexual and bisexual lizards different? Evidence from gastrointestinal parasites of *Darevskia* spp. in Turkey. *Acta Zool. Acad. Sci. Hung.*, 61(3): 279 – 288. DOI: 10.17109/AZH.61.3.6.2015
- ROCA, V., JORGE, F., ILGAZ, C., KUMLUTAS, Y., DURMUS, S.H., CARRETERO, M.A. (2016): Intestinal parasites of unisexual and bisexual lizards *Darevskia* spp. (Lacertidae) from Northeastern Anatolia. *Helminthologia*, 53(3): 298 – 303. DOI: 10.1515/helmin-2016-0021
- SAYGI, G., OLGUN, K. (1993): The nematode found in Dwarf Lizard (*Lacerta parva*) in Sivas province: *Spauligodon*. *Turk. Parazitol. Derg.*, 17(1): 40 – 45 (In Turkish)
- SCHAD, G.A., KUNTZ, R.E., WELLS, W.H. (1960): Nematode parasites from Turkish vertebrates. An annotated list. *Can. J. Zool.*, 38: 949 – 963. DOI: 10.1139/z60-101
- SINDACO, R., VENCHI, A., CARPANETO, G., BOLOGNA M. (2000): The reptiles of Anatolia. A checklist and zoogeographical analysis. *Biogeographia*, 21: 441 – 554
- Yamaguti, S. (1961): *Systema Helminthum*. Volume III. The Nematodes of Vertebrates, Part I. Interscience Publishers Inc., New York. 679 pp.
- YILDIRIMHAN, H.S., BURSEY, C.R., ALTUNEL, F.N. (2011): Helminth parasites of the Balkan green lizard *Lacerta trilineata* Bedriaga 1886 from Bursa, Turkey. *Turk. J. Zool.*, 35(4): 519 – 535. DOI: 10.3906/zoo-0910-1