



Fig. 224: *Eremias isfahanica* is characterized by scattered dark blotches on its flanks and two dark stripes without pale markings running parallel to the vertebral line. Behzad ZADHOUSH

**Status:** Common.

**Conservation status:** IUCN: Least Concern, CITES: –, National: Non-protected.

**Diet:** Arachnids and insects.

**Activity pattern:** Diurnal.

#### *Eremias isfahanica*

RASTEGAR-POUYANI, HOSSEINIAN, RAFIEE,

KAMI, RAJABIZADEH & WINK, 2016

Isfahan Racerunner

**Global range:** Iran.

**Distribution in Iran:** Isfahan Province.

**Biology:** The Isfahan Racerunner is found in wide desert regions with a scattered bushy vegetation. The habitat of this species are the plains with low, sparse vegetation beside mountains. Thus, inadvertent grazing of livestock is considered to be the main threatening factor for populations of this species.

**Size:** Snout–vent length 68 mm, tail length 114 mm.

**Reproduction:** Oviparous.





Fig. 225: *Eremias kavirensis*. Note white blotches on upper surface of legs, grey streaks with a brown margin and fingers edged with a row of comb-like scales. Kamran KAMALI

**Status:** Common.

**Conservation status:** IUCN: –, CITES: –, National: Non-protected.

**Diet:** Arachnids and insects.

**Activity pattern:** Diurnal.